

UC Irvine

UC Irvine Previously Published Works

Title

Convenient procedure for extraction of gonadotropin-releasing hormone and thyrotropin-releasing hormone from hypothalamic tissue.

Permalink

<https://escholarship.org/uc/item/38q6402t>

Journal

FEBS letters, 67(2)

ISSN

0014-5793

Authors

Koch, Y
Baram, T

Publication Date

1976-08-01

DOI

10.1016/0014-5793(76)80362-6

Copyright Information

This work is made available under the terms of a Creative Commons Attribution License, available at <https://creativecommons.org/licenses/by/4.0/>

Peer reviewed

CONVENIENT PROCEDURE FOR EXTRACTION OF GONADOTROPIN-RELEASING HORMONE AND THYROTROPIN-RELEASING HORMONE FROM HYPOTHALAMIC TISSUE

Y. KOCH and Tallie BARAM*

Department of Hormone Research, The Weizmann Institute of Science, Rehovot, Israel

Received 11 June 1976

1. Introduction

Measurement of tissue concentrations of the hypothalamic hormones gonadotropin-releasing hormone (GnRH) and thyrotropin-releasing hormone (TRH) is essential for studying their physiological roles, since reliable determination of the peripheral serum levels of these hormones has not yet been achieved [1,2]. Only minute amounts of GnRH and TRH are released into the hypothalamo-hypophyseal portal system, and these are greatly diluted in the general circulation. Moreover, these peptides have an extremely short half-life time [3,4]. Thus, at present, study of the fluctuations in the hypothalamic content of GnRH and TRH provides the only means of relating various physiological events to changes in the production and release of these hormones. The established procedures for quantitative extraction of hypothalamic hormones, however, are cumbersome as they involve acidification, readjustment of pH and solvent evaporation. This report presents an alternative, simple procedure for quantitative extraction of GnRH and TRH.

2. Materials and methods

2.1. Dissection

The medical basal hypothalamus (consisting mainly

of the median eminence and the arcuate nucleus; approximate weight 4–6 mg) was dissected, according to de Groot [5], from 5-month-old, Wistar-derived male rats of the departmental colony.

2.2. Extraction procedures

Hydrochloric acid: Hypothalamic tissue from one rat was homogenized in 1 ml of 0.1 N HCl and centrifuged at 14 000 g. The supernatant was neutralized (1 N NaOH), recentrifuged and used for radioimmunoassay.

Methanol or acetic-methanol: The tissue was homogenized in 1 ml methanol or in a solution of 2 N acetic acid in methanol. After centrifugation at 14 000 g, the solvent was removed by evaporation and the residue dissolved in 0.01 M phosphate-buffered saline (PBS; pH 6.9) for the assay.

Boiled neutral extract: The tissue was immersed in 1 ml PBS, in Thomas tissue grinder vessels (size AA), which were placed in a boiling bath for 3 min; homogenization in the same medium was followed by reboiling for 3 min and centrifugation at 14 000 g for 10 min. The supernatant fraction was kept for the assay.

2.3. Radioimmunoassay

Amounts of GnRH and TRH extracted by the four procedures described, were determined by radioimmunoassay, using antisera developed in this laboratory [6,7]. Samples were assayed in duplicate and at several dilutions. LH and TSH were assayed using kits kindly provided by the NIAMD-NIH Rat Pituitary Hormone Program. Results are expressed in terms of the RP-1 reference preparations.

* In partial fulfillment of the requirements for the Ph.D. degree of the Graduate School of the Weizmann Institute of Science.

2.4. Bioassay

Hemipituitaries of male rats were incubated for 2 h in 1 ml of Krebs-Ringer bicarbonate buffer containing 10 mg% glucose (KRBG). After change of medium the appropriate stimulants were added and incubations continued for 4 h. Amounts of LH and of TSH in the media were determined by radioimmunoassay.

3. Results

The concentrations of GnRH and TRH in rat hypothalamus, as determined by radioimmunoassay following the various extraction procedures, are summarized in table 1. Use of boiled neutral-PBS extracts yielded estimates of GnRH and TRH comparable to those resulting from the standard extraction procedures. The biological activity of GnRH and of TRH extracted by boiling, as compared to that of equivalent amounts of the corresponding synthetic peptides is shown in table 2. The extracted hormones are as potent in releasing LH or TSH as their synthetic counterparts.

4. Discussion

Data have been presented to show that GnRH and TRH can be quantitatively extracted by boiling tissue homogenates in an aqueous neutral solution;

Table 1
GnRH and TRH concentrations in the medial basal hypothalamus (MBH) of male rats, as estimated by use of four different extraction methods and an identical radioimmunoassay procedure

Extraction	GnRH ng/MBH	TRH ng/MBH
Boiled PBS	6.10 ± 0.26	4.86 ± 0.15
Methanol	5.81 ± 0.34	4.68 ± 0.23
Acetic-methanol	5.90 ± 0.36	4.45 ± 0.30
Hydrochloric acid	5.88 ± 0.37	4.48 ± 0.11

Mean ± S.E.M. for 12 tissue samples per group.

both hormones fully retain their biological and immunological activities. This short and simple technique makes it possible to use the extraction medium directly for radioimmunoassay or bioassay, eliminating the need for further manipulations such as neutralization or evaporation. Omitting the boiling step leads to poor yields, e.g.: less than 10% for GnRH, as observed by Shin and Howitt [8]. This finding may be explained by enzymic degradation of the peptides [9,10] and/or by their inaccessibility: the hormones may be bound to a protein carrier [11] or confined in vesicles [12]. Boiling inactivates degrading enzymes and may disrupt membrane structures or non-covalent protein-peptide binding. The biological and immunological properties

Table 2
Biological activity of hypothalamic GnRH and TRH extracted by boiling

Preparation	Hormone released to the medium (µg/ml)	
	LH	TSH
Control	0.59 ± 0.08 ^a	13.0 ± 1.7
Synthetic GnRH, 1 ng	5.10 ± 0.46	--
Hypothalamic GnRH, 1 ng ^b	5.53 ± 0.32	--
Synthetic TRH, 1 ng	--	138.0 ± 9.3
Hypothalamic TRH, 1 ng ^b	--	119.0 ± 9.8

^a Mean and standard error of the mean for 6–12 determinations.

^b As quantitated by radioimmunoassay.

Hypothalamic peptides were incubated for 4 h with hemipituitaries in 1 ml of KRBG. LH and TSH were determined by radioimmunoassay.

of the hypothalamic hormones, however, are unaffected by boiling.

Acknowledgements

We are grateful to Miss N. Ben Aroya for excellent technical assistance and to Dr H. R. Lindner for his interest in this study and assistance in the preparation of this manuscript. This work was supported by a grant (to H. R. Lindner) of the Ford Foundation and the Population Council Inc., N.Y.

References

- [1] Clemens, L. E., Kelch, R. P., Markovs, M., Westhoff, M. H. and Dermody, W. C. (1975) *J. Clin. Endocrinol. Metab.* 41, 1058–1064.
- [2] de la Cruz, K. G. and Arimura, A. (1975) In: 57th Annu. Meet. Endocr. Soc., New York, Abst. 106.
- [3] Redding, T. W. and Schally, A. V. (1972) *Neuroendocrinology* 9, 250–256.
- [4] Keye, W. R., Kelch, R. P., Niswender, G. D. and Jaffe, R. B. (1973) *J. Clin. Endocrinol. Metab.* 36, 1263–1267.
- [5] de Groot, J. (1972) in: *The rat forebrain in stereotaxic coordinates*, North-Holland Publishing Company, Amsterdam.
- [6] Koch, Y., Wilcheck, M., Fridkin, M., Chobsieng, P., Zor, U. and Lindner, H. R. (1973) *Biochem. Biophys. Res. Commun.* 55, 616–622.
- [7] Koch, Y., Baram, T. and Fridkin, M. (1976) *FEBS Lett.* 63, 295–299.
- [8] Shin, S. H. and Howitt, C. J. (1974) *Can. J. Physiol. Pharmacol.* 52, 754–758.
- [9] Lippman, F. (1973) *Acta Endocrin. Supp.* 180, 294–300.
- [10] Koch, Y., Baram, T., Chobsieng, P. and Fridkin, M. (1974) *Biochem. Biophys. Res. Commun.* 61, 95–103.
- [11] Fawcett, C. P. and Shin, S. H. (1975) *Endocrine Res. Commun.* 2, 151–158.
- [12] Ishii, S. (1970) *Endocrinology* 86, 207–216.